

FREE $\mathbf{C}_+$ -ACTIONS ON $\mathbf{C}^3$ ARE TRANSLATIONS

SHULIM KALIMAN¹

Department of Mathematics, University of Miami
Coral Gables, Florida 33124, USA
(e-mail: kaliman@math.miami.edu)

ABSTRACT. Let X' be a smooth contractible three-dimensional affine algebraic variety with a free algebraic $\mathbf{C}_+$ -action on it such that $S = X'//\mathbf{C}_+$ is smooth. We prove that X' is isomorphic to $S \times \mathbf{C}$ and the action is induced by a translation on the second factor. As a consequence we show that any free algebraic $\mathbf{C}_+$ -action on $\mathbf{C}^3$ is a translation in a suitable coordinate system.

1. Introduction. In 1968 Rentschler [Re] proved that every algebraic action of the additive group $\mathbf{C}_+$ of complex numbers on $\mathbf{C}^2$ is triangular in a suitable polynomial coordinate system. This implies that a free $\mathbf{C}_+$ -action on $\mathbf{C}^2$ (i.e. an action without fixed points) can be viewed as a translation. In 1984 Bass [Ba] found a $\mathbf{C}_+$ -action on $\mathbf{C}^3$ which is not triangular in any coordinate system. But the question about free $\mathbf{C}_+$ -actions on $\mathbf{C}^3$ remained open (e.g., see [Sn], [Kr], [DaFr], [DeFi]). While working on this problem we consider a more general situation when there is a nontrivial algebraic $\mathbf{C}_+$ -action on a complex three-dimensional affine algebraic variety X' such that its ring of regular functions is factorial and $H_3(X') = 0$. By a theorem of Zariski [Za] the algebraic quotient $X'//\mathbf{C}_+$ is isomorphic to an affine surface S . Let $\pi : X' \rightarrow S$ be the natural projection. Then there is a curve $\Gamma \subset S$ such that for $E = \pi^{-1}(\Gamma)$ the variety $X' \setminus E$ is isomorphic to $(S \setminus \Gamma) \times \mathbf{C}$ over $S \setminus \Gamma$. The study of morphism $\pi|_E : E \rightarrow \Gamma$ is central for this paper. As an easy consequence of the Stein factorization one can show that $\pi|_E = \theta \circ \vartheta$ where $\vartheta : E \rightarrow Z$ is a morphism into a curve Z with general fibers isomorphic to $\mathbf{C}$, and $\theta : Z \rightarrow \Gamma$ is a quasi-finite morphism. A more delicate fact (Proposition 3.2) is that θ is, actually, finite and, furthermore, if in addition X' is smooth and $H_2(X') = 0$, then each irreducible component Z^1 of Z , such that $\theta|_{Z^1}$ is

¹ The author was partially supported by the NSA grant MDA904-00-1-0016.

not injective, is rational and $\theta(Z^1)$ is a polynomial curve. A geometrical observation (Proposition 4.2) shows that in the case of a free $\mathbf{C}_+$ -action and a smooth S such a component Z^1 cannot exist. If the restriction of θ to any irreducible component of Z is injective then Γ can be chosen empty (Theorem 5.4), i.e. X' is isomorphic to $S \times \mathbf{C}$ over S . In combination with Miyanishi's theorem [Miy80] this yields the long-expected result.

Theorem 1. *Every free algebraic $\mathbf{C}_+$ -action on $\mathbf{C}^3$ is a translation in a suitable polynomial coordinate system.*

Acknowledgments. It is our pleasure to thank P. Bonnet, D. Daigle, G. Freudenburg, K.-H. Fieseler, and M. Zaidenberg for useful consultations.

2. Decomposition $\pi|_E = \theta \circ \vartheta$.

Lemma 2.1. *Let X' be a factorial affine algebraic variety of dimension 3 with a nontrivial $\mathbf{C}_+$ -action on it and $\pi : X' \rightarrow S$ be the quotient morphism. Then S is a factorial affine algebraic surface, $\pi^{-1}(s_0)$ is either a curve or empty for any $s_0 \in S$, and $E = \pi^{-1}(\Gamma)$ is a nonempty irreducible surface for every closed irreducible curve $\Gamma \subset S$. Furthermore, if $\Gamma = g^*(0)$ for a regular function g on S then $E = (g \circ \pi)^*(0)$.*

Proof. By [Za] S is an affine algebraic surface. Suppose that E is the union of a surface E' and a curve C where E' and C may be empty. Assume $\pi(E')$ is not dense in Γ . Consider a rational function h on S with poles on Γ only. Let e be the product of h and a regular function that does not vanish at general points of Γ but vanishes on $\pi(E')$ with sufficiently large multiplicity such that $e \circ \pi$ vanishes on $E' \setminus C$. As X' is factorial, by deleting singularities in codimension 2 we see that $e \circ \pi$ is a regular function on X' . As it is invariant under the action, e must be regular. Contradiction. Thus $\pi(E')$ is dense in Γ . Since X' is factorial E' is the zero fiber of a regular function f on X' . As f does not vanish on a general fiber $\pi^{-1}(s) \simeq \mathbf{C}$ it is constant on each general fiber. Hence f is invariant under the action, i.e. $f = g \circ \pi$ where g a regular function on S which implies that g is a defining function for Γ (i.e. S is factorial) and, furthermore, $E = E' \neq \emptyset$. If for a regular function f_1 on X' its zero fiber is an irreducible component of E then f_1 is again invariant under the action, i.e. $f_1 = g_1 \circ \pi$. As g_1 vanishes on Γ only it must be proportional to g . Thus E is irreducible and we get the last statement as well. If we assume that $\pi^{-1}(s_0)$ is a

surface for $s_0 \in S$ the similar argument yields a regular function on S that vanishes at a finite set only which is impossible. $\square$

If we have a nontrivial $\mathbf{C}_+$ -action on a three-dimensional affine algebraic variety X' with a quotient morphism $\pi : X' \rightarrow S = X'/\mathbf{C}_+$ then it is well-known that there is a closed curve $\Gamma \subset S$ such that for $\check{S} = S \setminus \Gamma$ and $E = \pi^{-1}(\Gamma)$ the variety $X' \setminus E = \pi^{-1}(\check{S})$ is naturally isomorphic to $\check{S} \times \mathbf{C}$ where the projection to the first factor corresponds to π . Let $\sigma_0 : X' \setminus E \rightarrow \check{S} \times \mathbf{C}$ be an isomorphism over $\check{S}$, i.e. $\sigma_0 = (\pi|_{X' \setminus E}, h_0)$ where h_0 is a regular function on $X' \setminus E$. There is some freedom in the choice of h_0 and, therefore, σ_0 .

Lemma 2.2. *Let X', E, Γ, S , and σ_0 be as before, $\Gamma = g^{-1}(0)$ for a regular function g on S , and $X = S \times \mathbf{C}$. Then σ_0 can be chosen so that there exists an affine modification $\sigma : X' \rightarrow X$ such that $\sigma|_{X' \setminus E} = \sigma_0$.*

Proof. Note that σ_0 extends to a rational map $\sigma = (\pi, h) : X' \rightarrow X$ where h is a rational function on X' whose restriction to $X' \setminus E$ coincides with h_0 . One can replace h_0 by $h_0 g^n$ (where we treat g as a function on X') as σ_0 remains an isomorphism after this replacement. Choosing $n > 0$ sufficiently large we can make h regular on X' whence σ becomes a regular birational morphism. $\square$

Lemma 2.3. *Let X' be a factorial affine algebraic variety of dimension 3 with a nontrivial $\mathbf{C}_+$ -action on it, $\pi : X' \rightarrow S$ be the quotient morphism, E and Γ be as before Lemma 2.2.*

(1) *There exist a curve Z , a morphism $\vartheta : E \rightarrow Z$ with connected general fibers, and a quasi-finite morphism $\theta : Z \rightarrow \Gamma$ for which $\pi|_E = \theta \circ \vartheta$. Furthermore, $\theta^{-1}(\Gamma^1)$ is irreducible for every irreducible component Γ^1 of Γ .*

(2) *There exists a finite set $F \subset \Gamma$ such that for $\Gamma^* = \Gamma \setminus F$, $Z^* = \theta^{-1}(\Gamma^*)$, and $E^* = \pi^{-1}(\Gamma^*)$ there is an isomorphism $E^* \simeq Z^* \times \mathbf{C}$ such that under this isomorphism $\vartheta|_{E^*}$ coincides with the projection to the first factor.*

Proof. Extend $\pi|_E : E \rightarrow \Gamma$ to a proper morphism $\bar{E} \rightarrow \bar{\Gamma}$ of complete varieties. By the Stein factorization theorem this extended morphism is a composition of morphisms $\bar{E} \rightarrow \bar{Z}$ and $\bar{Z} \rightarrow \bar{\Gamma}$ where the first morphism has connected fibers, $\bar{Z}$ is a curve, and the last morphism is finite. The restriction of these morphisms is what we need in (1). Since $E^1 = \pi^{-1}(\Gamma^1)$ is irreducible by Lemma 2.1, we see

that $\theta^{-1}(\Gamma^1) = \vartheta(E^1)$ is also irreducible. As E is the exceptional divisor of affine modification σ from Lemma 2.2, (2) follows from [Ka02, Cor. 2.2] (in the case of a free $\mathbf{C}_+$ -action it follows also from the fact that the general fibers of ϑ are orbits of the action, and, therefore, isomorphic to $\mathbf{C}$). $\square$

Remark 2.4. (1) Let the center C of the affine modification σ from Lemma 2.2 (i.e. C is the closure of $\sigma(E)$) be a curve. Note that $\theta : Z \rightarrow \Gamma$ factors through quasi-finite morphisms $Z \rightarrow C$ and $C \rightarrow \Gamma$. (2) Choosing “bigger” F and Γ one can always suppose that Γ^* , Z^* , and $S \setminus F$ are smooth.

3. Finiteness of θ .

Let S and F be as in Lemma 2.3. In the case of $S \simeq \mathbf{C}^2$ there is an isomorphism $H_0(F) \simeq H_3(S \setminus F)$: namely we assign to each $x_0 \in F$ the 3-cycle presented by a small sphere with center at x_0 . The same remains true in the normal case.

Lemma 3.1. *Let S be a normal affine surface, Γ be a closed curve in S , $\check{S} = S \setminus \Gamma$, F be a finite subset of Γ such that $\Gamma^* = \Gamma \setminus F$ and $\tilde{S} = S \setminus F$ are smooth.*

(i) *There exist a natural isomorphism $\phi : H_0(F) \rightarrow H_3(\tilde{S})$.*

(ii) *Let $\varphi : H_3(\tilde{S}) \rightarrow H_1(\Gamma^*)$ be the composition of the Thom isomorphism $H_3(\tilde{S}, \check{S}) \rightarrow H_1(\Gamma^*)$ and the natural homomorphism $H_3(\tilde{S}) \rightarrow H_3(\tilde{S}, \check{S})$, let $x_0 \in F$, $\omega = \phi(x_0)$, and Γ_j , $j = 1, \dots, k$ be irreducible analytic branches of Γ at x_0 . Suppose that γ_j is a simple loop in Γ_j around x_0 with positive orientation. Then $\varphi(\omega) = \sum_{j=1}^k \gamma_j$ (where we treat γ_j as an element of $H_1(\Gamma^*)$).*

Proof. As $H_3(S) = H_4(S) = 0$ [Mil, Th. 7.1] the exact homology sequence of pair $(S, \tilde{S})$ shows that $H_3(\tilde{S}) \simeq H_4(S, \tilde{S})$. Thus in (i) it suffices to prove that $H_4(S, \tilde{S}) \simeq H_0(F)$. As S may not be smooth we cannot apply Thom’s isomorphism. But by the excision theorem $H_4(S, \tilde{S}) \simeq H_4(U, U \setminus F)$ where U is a small Euclidean neighborhood of F in S , i.e. U is a disjoint union $\bigcup_i U_i$ of neighborhoods U_i of a point x_i running over F . Hence $H_4(S, \tilde{S}) = \oplus_i H_4(U_i, U_i \setminus x_i)$. Let B_ϵ be the ball of radius ϵ in $\mathbf{C}^n \supset S$ with center at $x_0 \in F$ and $U_0 = S \cap B_\epsilon$. For small $\epsilon > 0$, $\omega = S \cap \partial B_\epsilon$ is a three-dimensional real manifold and U_0 is a cone over ω [Lo]. Hence $H_4(U_0, U_0 \setminus x_0) \simeq H_3(U_0 \setminus x_0) \simeq \mathbf{Z}$ and U_0 and ω are generators of $H_4(U_0, U_0 \setminus x_0)$ and $H_3(U_0 \setminus x_0)$ respectively. Thus $H_4(S, \tilde{S}) \simeq H_0(F)$ which is (i). For small $\epsilon > 0$ the sphere ∂B_ϵ and, thus, ω meet each Γ_j transversally along a simple loop γ_j around x_0

in Γ_j . Hence the description of Thom's isomorphism [Do] yields $\varphi(\omega) = \sum_{j=1}^k \gamma_j$. $\square$

Proposition 3.2. *Let the assumption and notation of Lemma 2.3 hold, $H_3(X') = 0$, and $\tilde{X} = \pi^{-1}(\tilde{S})$ be smooth where $\tilde{S} = S \setminus F$.*

(i) *Then θ is a finite morphism.*

(ii) *If x_0 is a selfintersection point of Γ then $\theta^{-1}(x_0)$ consists of one point z_0 .*

(iii) *Let X' be also smooth and $H_2(X') = 0$. Let Z^1 be an irreducible component of Z and $\Gamma^1 = \theta(Z^1)$. If $\theta|_{Z^1} : Z^1 \rightarrow \Gamma^1$ is not bijective then Γ^1 is a polynomial curve (i.e. its normalization is $\mathbf{C}$) and Z^1 is rational.*

Proof. Put $\check{X} = X' \setminus E$ and $\check{S} = S \setminus \Gamma$, i.e. $\check{X} \simeq \check{S} \times \mathbf{C}$. We have the following commutative diagram of the exact homology sequences of pairs.

$$\begin{array}{ccccccccc}
\cdots & \longrightarrow & H_{j+1}(\tilde{X}, \check{X}) & \longrightarrow & H_j(\check{X}) & \longrightarrow & H_j(\tilde{X}) & \longrightarrow & H_j(\tilde{X}, \check{X}) & \longrightarrow & H_{j-1}(\check{X}) & \longrightarrow \cdots \\
& & \downarrow & & \simeq \downarrow & & \downarrow & & \downarrow & & \simeq \downarrow & \\
\cdots & \longrightarrow & H_{j+1}(\tilde{S}, \check{S}) & \longrightarrow & H_j(\check{S}) & \longrightarrow & H_j(\tilde{S}) & \longrightarrow & H_j(\tilde{S}, \check{S}) & \longrightarrow & H_{j-1}(\check{S}) & \longrightarrow \cdots
\end{array}$$

which we consider for $j = 3$. By Remark 2.4 we can suppose that Γ^* , $\tilde{S}$, and E^* are smooth. As $\tilde{X} \setminus \check{X} = E^*$ the Thom isomorphism [Do] implies $H_i(\tilde{X}, \check{X}) \simeq H_{i-2}(E^*)$ and, similarly, $H_i(\tilde{S}, \check{S}) \simeq H_{i-2}(\Gamma^*)$. As $H_3(\check{S}) = 0$ [Mil, Th. 7.1] and $H_i(E^*) = H_i(Z^*)$ we have

$$\begin{array}{ccccccccc}
\cdots & \longrightarrow & 0 & \longrightarrow & H_3(\tilde{X}) & \xrightarrow{\varphi'} & H_1(Z^*) & \xrightarrow{\psi'} & H_2(\check{X}) & \xrightarrow{\chi'} & H_2(\tilde{X}) & \longrightarrow \cdots \\
& & \downarrow & & \downarrow \delta_3 & & \downarrow \theta_1 & & \simeq \downarrow \check{\delta}_2 & & \downarrow \delta_2 & \\
\cdots & \longrightarrow & 0 & \longrightarrow & H_3(\tilde{S}) & \xrightarrow{\varphi} & H_1(\Gamma^*) & \xrightarrow{\psi} & H_2(\check{S}) & \xrightarrow{\chi} & H_2(\tilde{S}) & \longrightarrow \cdots
\end{array} \tag{*}$$

As $H_4(X') = 0$ [Mil, Th. 7.1] the exact homology sequence of pair $(X', \tilde{X})$ implies that $H_3(\tilde{X})$ is isomorphic to $H_4(X', \tilde{X})$. By Lemma 2.1 $L = \pi^{-1}(F) = E \setminus E^* = X' \setminus \tilde{X}$ is a curve. If U is a small Euclidean neighborhood of L in X' then $H_4(U, U \setminus L) = H_4(X', \tilde{X})$ by the excision theorem. That is, every 3-cycle $\omega' \in H_3(\tilde{X})$ can be chosen in $U \setminus L$. Thom's isomorphism maps the image of ω' in $H(U, U \setminus L)$ to a 1-cycle $\alpha \in H_1(E^* \cap U)$. As Thom's isomorphisms are functorial under open embeddings [Do, Ch. 8, 11.5] we can treat α as an element of $H_1(E^*)$. Thus the image of α under

the homomorphism generated by the natural projection $E^* \rightarrow Z^*$ coincides with $\beta_0 = \varphi'(\omega') \in H_1(Z^*)$. As α is contained in a small neighborhood $U \cap E$ of L in E we see that β_0 is contained in a small neighborhood of the finite set $Z \setminus Z^*$. This implies that $\text{Im } \varphi'$ is contained in the kernel G' of the homomorphism $H_1(Z^*) \rightarrow H_1(Z)$ induced by the natural embedding $Z^* \hookrightarrow Z$. Let $\bar{\theta} : \bar{Z} \rightarrow \bar{\Gamma}$ be the extension of θ to completions of Z and Γ . Assume that $F' = (\bar{Z} \setminus Z) \setminus \bar{\theta}^{-1}(\bar{\Gamma} \setminus \Gamma)$ is not empty which is equivalent to the fact that θ is not finite. Let $z_0 \in F'$. One can suppose that $x_0 = \theta(z_0) \in F$. Let Γ_j , $j = 1, \dots, k$ be irreducible analytic branches of Γ at x_0 , and let γ_j be a simple loop in Γ_j around x_0 with positive orientation. We treat $\gamma := \sum_{j=1}^k \gamma_j$ as an element of $H_1(\Gamma^*)$. By Lemma 3.1 $H_3(\tilde{S})$ is generated by elements ω of form $\omega = \phi(x_0)$ and $\varphi(\omega) = \gamma$. Take a simple loop β_1 around z_0 in an irreducible analytic branch Z_1 of $\bar{Z}$ at z_0 such that $\bar{\theta}(Z_1)$ is, say, Γ_1 . Take other simple loops around the points of $\bar{\theta}^{-1}(x_0)$ whose images under θ are contained in $\bigcup_{j=2}^k \Gamma_j$. Then we can construct $\beta \notin G'$ as an integer linear combination of β_1 and these other loops so that $\theta_1(\beta) = m\gamma$ where $m > 0$. As $\gamma \in \text{Im } \varphi$ we have $\psi(\gamma) = 0$. As $\check{\delta}_2$ is an isomorphism we have $\psi'(\beta) = 0$, i.e. $\beta \in \text{Im } \varphi'$. This contradiction implies (i).

For (ii) we need another look at $\text{Im } \varphi'$. Let $Z \setminus Z^* = \{z_i\}$ and G'_i be the kernel of the homomorphism $H_1(Z \setminus z_i) \rightarrow H_1(Z)$ induced by the natural embedding $Z \setminus z_i \hookrightarrow Z$. Note that G' is naturally isomorphic to $\bigoplus_i G'_i$. Moreover, since L is the disjoint union $\bigcup_i \vartheta^{-1}(z_i)$ one can choose a neighborhood U of L in X' as a disjoint union of neighborhoods of the curves $\vartheta^{-1}(z_i)$. Hence, repeating the argument with the excision theorem, we get $\text{Im } \varphi' = \bigoplus_i (G'_i \cap \text{Im } \varphi')$. Let x_0 be a selfintersection point of Γ (i.e. the number k of irreducible analytic branches Γ_j of Γ at x_0 is at least 2) and assume for simplicity that $\theta^{-1}(x_0)$ consists of two points z_0 and z_1 . Note that $z_0, z_1 \in Z \setminus Z^*$. Let $\beta' \in G'_0 \cap \text{Im } \varphi'$ (i.e. $\beta' = \varphi'(\omega')$). Then $\delta_3(\omega') = m\omega$ where $m \in \mathbf{Z}$ and ω is as before. As $\varphi(m\omega) = \theta_1(\beta')$ we see that $\theta_1(\beta') = m\gamma$. Similarly $\theta_1(\beta'')$ is proportional to γ for every $\beta'' \in G'_1 \cap \text{Im } \varphi'$. On the other hand since x_0 is a selfintersection point, we can find $\gamma' \in G'_0$ and $\gamma'' \in G'_1$ such that $\theta_1(\gamma')$ and $\theta_1(\gamma'')$ are not proportional to γ but $\theta_1(\gamma' + \gamma'')$ is. The commutativity of (*) implies that $\gamma' + \gamma'' \in \text{Ker } \psi' = \text{Im } \varphi'$. But the decomposition of $\text{Im } \varphi'$ we discussed before, implies that γ' and γ'' must be in $\text{Im } \varphi'$. This contradiction implies (ii).

Let G be the kernel of the homomorphism $H_1(\Gamma^*) \rightarrow H_1(\Gamma)$, induced by the natural embedding. There is a non-canonical isomorphism $H_1(\Gamma^*) = G \oplus H_1(\Gamma^{\text{norm}})$

(resp. $H_1(Z^*) = G' \oplus H_1(Z^{\text{norm}})$) such that $\text{Im } \varphi$ (resp. $\text{Im } \varphi'$) is contained in the first term and $\theta_1(G')$ is a subgroup of finite index in G . As $\check{\delta}_2$ is an isomorphism the ranks of $H_1(\Gamma^{\text{norm}})$ and $H_1(Z^{\text{norm}})$ are the same, and the ranks of $G/\text{Im } \varphi$ and $G'/\text{Im } \varphi'$ coincide as well. For (iii) note that (i) and (ii) implies that $\theta|_{Z^1} : Z^1 \rightarrow \Gamma^1$ is bijective as soon as it is birational. If it is not birational then the fact, that the ranks of $H_1(\Gamma^{\text{norm}})$ and $H_1(Z^{\text{norm}})$ are the same, implies that the genus of Z^1 (and, therefore, Γ^1) is zero. Assume that the normalization of Γ^1 is different from $\mathbf{C}$, i.e. one can suppose that $\bar{\Gamma}^1 \setminus \Gamma^1$ consists of more than one point where $\bar{\Gamma}^1$ is the closure of Γ^1 in $\bar{\Gamma}$. Let $\bar{Z}^1$ be the closure of Z^1 in $\bar{Z}$. Then either $\bar{\theta}$ maps $\bar{Z}^1 \setminus Z^1$ bijectively onto $\bar{\Gamma}^1 \setminus \Gamma^1$ or it does not. In the first case a simple loop in Γ^1 around a point $x \in \bar{\Gamma}^1 \setminus \Gamma^1$ produces $\gamma^1 \in H_1(\Gamma^*)$ which is not in $G + \text{Im } \theta_1$ and in the second case an integer linear combination of simple loops in Z^1 around points of $\bar{\theta}^{-1}(x)$ produces $\beta^1 \in H_1(Z^*)$ which is in $\text{Ker } \theta_1 \setminus G'$. In the first case since $G \supset \text{Im } \varphi = \text{Ker } \psi$, we see that $\varphi(\gamma^1)$ is not in $\text{Im } \check{\delta}_2 \circ \psi'$. On the other hand, as X' is smooth and $H_2(X') = 0$ we get $H_2(\tilde{X}) = 0$ whence $\text{Im } \psi' = H_2(\tilde{X})$. This is a contradiction since $\check{\delta}_2$ is an isomorphism. The second case is similar which concludes (iii). $\square$

Remark 3.3. Proposition 3.2 implies that π is surjective which is a generalization of the result of Bonnet [Bo] who proved this in the case of $X' \simeq \mathbf{C}^3$. This implies that when X' is Cohen-Macaulay, the algebra of regular functions on X' is faithfully flat over the algebra of regular functions on S (this follows from [Ma, Chap. 2, (3.J)] and Th. 3] and [Ei, Th. 18.16], see also [Da]).

4. Injectivity of θ .

Lemma 4.1. *Let the notation of Lemma 2.3 hold, $\mathcal{M} \subset E$ be a germ of a curve at a general point x' of E such that $\pi|_{\mathcal{M}} : \mathcal{M} \rightarrow S$ is injective. Let $\mathcal{P} \subset X'$ be a germ of a surface at x' that meets E transversally along $\mathcal{M}$. Then $\pi|_{\mathcal{P}} : \mathcal{P} \rightarrow S$ is injective.*

Proof. While speaking of affine modifications here and in Section 5 we use the terminology from [Ka02]. Consider the affine modification $\sigma : X' \rightarrow X \simeq S \times \mathbf{C}$ from Lemma 2.2. Proposition 2.12 from [Ka02] implies that in a neighborhood of a general point of E this modification σ is a composition of basic modifications. Locally, these basic modifications are nothing but usual monoidal transformations [Ka02, Remark 2.3] (see also Remark 5.2 below). In the case of three-dimensional manifolds and one-dimensional centers the coordinate form of these monoidal transformations is

$(\xi_1, \xi_2, \xi_3) \rightarrow (\xi_1, \xi_1 \xi_2, \xi_3) := (\zeta_1, \zeta_2, \zeta_3)$. If σ is locally such a monoidal transformation, then σ maps $\mathcal{M}$ injectively into the line $\zeta_1 = \zeta_2 = 0$ since $\pi = \text{pr} \circ \sigma$ where $\text{pr} : X \rightarrow S$ is the natural projection. Hence $\mathcal{M}$ must be given by equations of form $\xi_1 = \xi_2 - g_0(\xi_3) = 0$. As $\mathcal{P}$ meets E transversally along $\mathcal{M}$ we see that $\mathcal{P}$ is given by an equation of form $\xi_2 - g(\xi_1, \xi_3) = 0$ where $g(0, \xi_3) = g_0(\xi_3)$. Hence the image of $\mathcal{P}$ is the germ of a surface given by $\zeta_2 = \zeta_1 g(\zeta_1, \zeta_3)$. In particular, this germ is isomorphic to $\mathcal{P}$ (via the transformation) and transversal to the plane $\zeta_1 = 0$. Induction on the number of monoidal transformations in the local decomposition of σ implies that $\sigma|_{\mathcal{P}} : \mathcal{P} \rightarrow \mathcal{Q} := \sigma(\mathcal{P})$ is bijective and $\mathcal{Q}$ meets D transversally along a germ $\mathcal{C}$ of $C = \sigma(E)$ at $x = \sigma(x')$. As x is a general point of C , projection pr maps $\mathcal{C}$ injectively into Γ . As $\mathcal{Q}$ meets D transversally along $\mathcal{C}$, morphism $\text{pr}|_{\mathcal{Q}} : \mathcal{Q} \rightarrow S$ is not ramified along $\mathcal{C}$ whence this map is injective. Thus $\pi|_{\mathcal{P}} : \mathcal{P} \rightarrow S$ is injective. $\square$

Proposition 4.2. *Let the assumption of Proposition 3.2 (iii) hold, S be smooth, and $\pi : X' \rightarrow S$ be the quotient morphism of a free $\mathbf{C}_+$ -action Φ on X' . Then θ is bijective.*

Proof. Assume the contrary, i.e. there is a component Z^1 of the Z such that $\theta|_{Z^1}$ is not bijective. By Proposition 3.2 (i)-(ii) $\theta|_{Z^1}$ is not birational. Furthermore, Proposition 3.2 (iii) implies that Z contains a ramification point z_0 of θ whose order is $m \geq 2$. Take a smooth point x'_0 of $\vartheta^{-1}(z_0)$. The irreducible component L_0 of $\vartheta^{-1}(z_0)$, that contains x'_0 , is an irreducible component of a fiber of π and, therefore, an orbit of Φ since Φ is free. Consider the germ $\mathcal{P}$ of a surface at x'_0 transversal to L_0 . Then all orbits of Φ close to L_0 must be transversal to $\mathcal{P}$. Let us show that this cannot be true.

Consider $\varrho = \pi|_{\mathcal{P}} : \mathcal{P} \rightarrow \pi(\mathcal{P})$. One can see that ϱ is finite (as $\mathcal{P}$ does not contain fibers of π). As $\pi|_E = \theta \circ \vartheta$ and θ has a ramification point of order m we see that ϱ is at least m -sheeted. As S is smooth there must be a ramification curve $R \subset \mathcal{P}$ of morphism ϱ . Consider a general point x'_1 of the curve $\mathcal{M} = \mathcal{P} \cap E$, and the germ $\mathcal{P}_1$ of $\mathcal{P}$ at x'_1 . Note that $\mathcal{P}_1$ meets E transversally along the irreducible germ $\mathcal{M}_1$ of $\mathcal{M}$ at x'_1 , and $\pi|_{\mathcal{M}_1}$ is injective. By Lemma 4.1, ϱ is locally injective in a neighborhood of x'_1 , i.e. R has no common components with $\mathcal{M}$. Therefore, a general point $x'_2 \in R$ is contained in $X' \setminus E \simeq \check{S} \times \mathbf{C}$. As x'_2 is a ramification point of ϱ it cannot be a point where $\mathcal{P}$ meets the fiber of the natural projection $X' \setminus E \rightarrow \check{S}$ (which is a fiber of π

and, therefore, an orbit of Φ) transversally which yields the desired contradiction. $\square$

5. Main theorem.

We want to decompose $\sigma : X' \rightarrow X$ from Lemma 2.2 into a composition of simpler modifications $X' \rightarrow X_1$ and $X_1 \rightarrow X$. Let A , A_1 , and A' be the algebras of regular functions on X , X_1 , and X' respectively. In order to find X_1 it suffices to find an affine domain A_1 which is a proper subring of A' and which contains but not equal to A .

Lemma 5.1. *Let $\sigma : X' \rightarrow X$ be an affine modification, C be its center, D^1 be an irreducible component of its divisor D , $C^1 = C \cap D^1$. Suppose that $A' = A[I/f]$ where I is an ideal in A that vanish on C , $f = g^n h$, $D = f^{-1}(0)$, and $D^1 = g^*(0)$ while h is not identically zero on D^1 . Let $E^1 = \sigma^{-1}(C^1)$, and the defining ideal $\mathcal{I}_{A'}(E^1)$ of E^1 be the principal ideal in A' generated by g .*

(1) *Then $A[(\mathcal{I}_A(C^1))^n/g^n] \subset A'$ for the defining ideal $\mathcal{I}_A(C^1)$ of C^1 in A . In particular, for every ideal $J \subset \mathcal{I}_A(C^1)$ the algebra $A_1 = A[J/g]$ is contained in A' .*

(2) *If J in (1) coincides with $\mathcal{I}_A(C^1)$ then $A' = A_1[I_1/f_1]$ where I_1 is an ideal in A_1 and $f_1 = g^{n-1}h$.*

Proof. Note that for every $a \in (\mathcal{I}_A(C^1))^n$ we have $a \in (\mathcal{I}_{A'}(E^1))^n$ (as an element of A') whence $a/g^n \in A'$. For the second statement of (1) note that $A_1 = A[(g^{n-1}J)/g^n] \subset A[(\mathcal{I}_A(C^1))^n/g^n]$.

Let $g, b_1, \dots, b_k$ be generators of $I \subset \mathcal{I}_A(C^1)$. Then in (2) $b_1/g, \dots, b_k/g \in A_1$. The ideal I_1 generated by these elements is what we need in (2). $\square$

Remark 5.2. Let $x \in C^1$ be a smooth point of both C^1 and X , and the zero multiplicity of g at x be 1. In a small Euclidean neighborhood U of X the defining ideal of $C^1 \cap U$ is a complete intersection, i.e. it is generated by regular functions $g, b_1, \dots, b_k$ which are elements of a local coordinate system on U . Hence in U the modification $A \hookrightarrow A_1 = A[\mathcal{I}_A(C^1)/g]$ can be viewed as a monoidal transformation. This fact in combination with Lemma 5.1 (2) and induction on n yields another way to show that modification σ in the proof of Lemma 4.1 is a composition of monoidal transformations in a neighborhood of a general point of its center.

Proposition 5.3. *Let the assumption of Proposition 3.2 (iii) hold, and Γ^1 (resp. Z^1) be an irreducible components of Γ (resp. Z) such that $\theta(Z^1) \subset \Gamma^1$. Suppose that*

$\theta|_{Z^1} : Z^1 \rightarrow \Gamma^1$ is bijective. Let $\Gamma^\bullet$ be the closure of $\Gamma \setminus \Gamma^1$ in S . Then $\pi^{-1}(S \setminus \Gamma^\bullet)$ is isomorphic to $(S \setminus \Gamma^\bullet) \times \mathbf{C}$ over $S \setminus \Gamma^\bullet$.

Proof. Let $\sigma : X' \rightarrow X \simeq_S S \times \mathbf{C}$ be the affine modification from Lemma 2.2. In particular, its divisor $D = \Gamma \times \mathbf{C}$ contains an irreducible component $D^1 = \Gamma^1 \times \mathbf{C}$. Our aim is to construct another affine modification $\sigma^\bullet : X' \rightarrow X$ such that its divisor is $D^\bullet = \Gamma^\bullet \times \mathbf{C}$ which would yield the statement of this Proposition. As S is factorial by Lemma 2.1, Γ^1 is the zero locus of a regular function g on S . If we treat g as a regular function on X we get $A' = A[I/f]$ where I and $f = g^n h$ are as in Lemma 5.1. In virtue of [Ka02, Lemma 2.2] one can suppose that the center C of σ is a curve. By Remark 2.4 $\theta : Z \rightarrow \Gamma$ factors through $\text{pr}|_C : C \rightarrow \Gamma$ and $Z \rightarrow C$. Let C^1 be the image of Z^1 under the last morphism. Then $\text{pr}|_{C^1} : C^1 \rightarrow \Gamma^1$ is bijective. Hence C^1 is given in $D^1 = \Gamma^1 \times \mathbf{C}$ by $t - \bar{b}_1 = 0$ where t is a coordinate on the second factor of $X \simeq S \times \mathbf{C}_t$ and $\bar{b}_1$ is a continuous rational function on Γ^1 . Let us show that $t - \bar{b}_1$ is regular on D^1 . Let x_0 be the preimage of $o \in \Gamma^1$ in C^1 and $x'_0 \in \pi^{-1}(o)$ be a smooth point of $\pi^{-1}(o)$. Take the germ $\mathcal{P}$ of an analytic surface at x'_0 such that $\mathcal{P} \cap L = x'_0$. One can check that $\mathcal{Q} = \sigma(\mathcal{P})$ is closed in a neighborhood of x_0 and meets D_1 along the germ $\mathcal{C}^1$ of C^1 at x_0 . Hence $\mathcal{Q}$ is given in $X \simeq S \times \mathbf{C}_t$ by zeros of a function $t^n + a_{n-1}t^{n-1} + \dots + a_0$ where each a_i is a germ of a continuous function at $o \in S$ which is analytic outside Γ^1 . As S is normal a_i is analytic at o as well. Hence by [Ka02, Lemma 3.3] $t - \bar{b}_1$ is regular. Let b_1 be an extension of $\bar{b}_1$ to S . Then $A_1 = A[(t - b_1)/g]$ is the algebra of regular functions on a hypersurface X_1 in $X \times \mathbf{C}_u$ given by equation $ug = t - b_1$, and $X_1 \simeq S \times \mathbf{C}_u \simeq X$. By Lemma 2.1 the defining ideal of $E^1 = \sigma^{-1}(C^1)$ in A' is the principal ideal generated by g . As Lemma 2.3 implies that $Z^1 = \theta^{-1}(\Gamma^1)$, we have $C^1 = D^1 \cap C$. Thus by Lemma 5.1 $A_1 \subset A'$. Hence we can replace modification $\sigma : X' \rightarrow X$ by modification $X' \rightarrow X_1 \simeq_S S \times \mathbf{C}$ (induced by the embedding $A_1 \hookrightarrow A'$) and all assumption of Proposition 3.2 hold but the advantage is that now $f = g^{n-1}h$ by Lemma 5.1 (since the defining ideal of C^1 in A is generated by g and $t - b_1$). Repeating this procedure we can make $n = 0$, i.e. we get the modification $\sigma^\bullet : X' \rightarrow X$ which concludes the proof. $\square$

Theorem 5.4. *Let the assumption of Lemma 2.3 hold, X' be smooth, and $H_2(X') = H_3(X') = 0$.*

(i) *Then Γ can be chosen so that each irreducible component Z^1 of Z (resp. $\Gamma^1 =$*

$\theta(Z^1)$ of Γ) is rational (resp. a polynomial curve), and $\theta|_{Z^1} : Z^1 \rightarrow \Gamma^1$ is not bijective.

(ii) If S is smooth and the $\mathbf{C}_+$ -action Φ on X' has no fixed points then X' is isomorphic to $S \times \mathbf{C}$ over S and the action is generated by a translation on the second factor.

Proof. Applying Proposition 5.2 consequently to each component Z^1 of Z , such that $\theta|_{Z^1} : Z^1 \rightarrow \Gamma^1 = \theta(Z^1)$ is bijective, we obtain (i) in virtue of Proposition 3.2 (iii). Proposition 4.2 and (i) imply that the quotient morphism $\pi : X' \rightarrow S$ makes X' isomorphic to $S \times \mathbf{C}$ over S . $\square$

Proof of Theorem 1. By [Miy80] $\mathbf{C}^3//\mathbf{C}_+ \simeq \mathbf{C}^2$ for every nontrivial $\mathbf{C}_+$ -action on $\mathbf{C}^3$ whence Theorem 5.4 (ii) implies the desired conclusion. $\square$

Remark 5.5. (1) There is another way to extract Theorem 1 from Proposition 4.2. For a bijective θ and $X' \simeq \mathbf{C}^3$ an easy Euler characteristics argument shows that all fibers of ϑ (not only the general ones) are irreducible. This, in turn, implies that the $\mathbf{C}_+$ -action is separated and, therefore, it is a translation [DeFi, Kr].

(2) Consider an n -dimensional smooth contractible affine algebraic variety X' and a free algebraic action of a unipotent group U on it (i.e. each orbit of the action is isomorphic to U). Suppose that U is of dimension $n - 2$ (i.e. U is isomorphic to $\mathbf{C}^{n-2}$ as an affine algebraic variety). By Fujita's result (e.g., see [Ka94, Prop. 3.2]) X' is factorial and by Zariski's theorem [Za] X'/U is still an affine algebraic surface whence one can repeat all Lemmas and Propositions before for this situation with obvious adjustments (say, in Lemma 2.3 E^* will be isomorphic to $Z^* \times \mathbf{C}^{n-2}$). As Miyanishi's theorem [Miy80] says that in the case of a free action $\mathbf{C}^n//U \simeq \mathbf{C}^2$, we have the following analogue of Theorem 1: for every free algebraic action of U on $\mathbf{C}^n$ there is an isomorphism $\mathbf{C}^n \simeq \mathbf{C}^2 \times U$ such that the action is induced by the natural action on the second factor.

REFERENCES

- [Ba] H. Bass, *A nontriangular action of G_a on A^3* , J. Pure Appl. Algebra **33** (1984), no. 1, 1–5.
- [Bo] P. Bonnet, *Surjectivity of quotient maps for algebraic $(\mathbf{C}, +)$ -actions and polynomial maps with contractible fibers*, Trans. Groups (to appear).

- [Da] D. Daigle, *A note on generic planes and locally nilpotent derivations of $k[X, Y, Z]$* , (to appear).
- [DaFr] D. Daigle, G. Freudenburg, *Locally nilpotent derivations over a UFD and an application to rank two locally nilpotent derivations of $k[X_1, \dots, X_n]$* , J. Algebra **204** (1998), no. 2, 353–371.
- [DeFi] J. Deveney, D. Finston, *Free G_a actions on $\mathbf{C}^3$* , Proc. Amer. Math. Soc. **128** (2000), no. 1, 31–38.
- [Do] A. Dold, *Lectures on Algebraic Topology*, Springer, Berlin e.a., 1974.
- [Ei] D. Eisenbud, *Commutative Algebra with a View Towards Algebraic Geometry*, Graduate Texts in Math., Springer, N.Y. e.a., 1994.
- [Ka94] S. Kaliman, *Exotic analytic structures and Eisenman intrinsic measures*, Israel Math. J. **88** (1994), 411–423.
- [Ka02] S. Kaliman, *Polynomials with general $\mathbf{C}^2$ -fibers are variables*, Pacific J. Math., **203** (2002), no. 1, p. 161–189.
- [Kr] H. Kraft, *Challenging problems on affine n -space*, Séminaire Bourbaki, Vol. 1994/95; Astérisque **237** (1996), 295–317.
- [Lo] E.J.N. Looijenga, *Isolated singular points on complete intersections*, London Mathematical Society Lecture Note Series, **77**, Cambridge University Press, Cambridge, 1984, 200 p.
- [Ma] H. Matsumura, *Commutative Algebra*, The Benjamin/Cummings Publishing Company, Reading, Massachusetts, 1980.
- [Mil] J. Milnor, *Morse Theory*, Princeton Univ. Press, Princeton, NJ, 1963.
- [Miy80] M. Miyanishi, *Regular subrings of a polynomial ring*, Osaka J. Math. **17** (1980), no. 2, 329–338.
- [Re] R. Rentschler, *Opérations du groupe additif sur le plan affine*, C.R. Acad. Sci. Paris, **267** (1968), 384–387.
- [Sn] D. Snow, *Unipotent actions on affine space*, in “Topological methods in algebraic transformation groups”, Birkhäuser, Boston, MA, 1989, 165–176.
- [Za] O. Zariski, *Interpretations algébriques-géométriques du quatorzième problème de Hilbert*, in “O. Zariski: Collected Papers”, **2**, The MIT Press, Cambridge (Massachusetts), 1973, 261–275.